

THE EXTENSION OF BUCKLEY-FEURING SOLUTIONS FOR NON-POLYNOMIAL FUZZY PARTIAL DIFFERENTIAL EQUATIONS

D. GÁLVEZ AND J. L. PINO

DEPARTAMENTO DE ESTADÍSTICA E INVESTIGACIÓN OPERATIVA
UNIVERSIDAD DE SEVILLA

ABSTRACT. This paper presents the natural extension of Buckley-Feuring method proposed in [2] for solving fuzzy partial differential equations (FPDE) in a non-polynomial relation, such as the operator $\varphi(D_{x_1}, D_{x_2})$, which maps to the quotient between both partials. The new assumptions and conditions proceedings from this consideration are given in this document.

INTRODUCTION

Many approaches for obtaining non-numerical solutions of fuzzy differential equations (FDE) have been developed from the introduction of fuzzy set concept by Zadeh [7]. These ones give a diversity of definitions for FDE solution based on different notions of fuzzy derivative, such as Seikkala derivative, Buckley-Feuring derivative, Puri-Ralescu derivative, Kandel-Friedman-Ming derivative, Goetschel-Voxman derivative, or Dubois-Prade derivative. Some relations between these derivatives are presented by Buckley and Feuring in [3]. However, only a few of these fuzzy derivatives are valid in some contexts as FDE solution. For example, the Goetschel-Voxman derivative, or the Dubois-Prade derivative provide solutions that cannot be a fuzzy number, whereas Puri-Ralescu derivative, and Kandel-Friedman-Ming derivative, always exist and provide a fuzzy number as solution of the FDE, but making use of abstract subtractions of fuzzy concepts in their definitions, making difficult the interpretation of this solutions in some real applications.

This paper uses the Buckley-Feuring derivative for solving FPDE. This derivative does not always exist, but if it does, provides a fuzzy number solution easily understandable in the context in which a specific FPDE has been developed.

The authors which proposed this concept of derivative, developed a methodology for solving constant coefficients polynomial FPDE in [2]. This paper present the extension of this methodology to a non-polynomial expression in partial fuzzy derivatives.

In the following lines, the components of a FPDE are enumerated:

- x_i , $i = 1, 2$, $x_1 \in S_1 \subset I_1 = (0, M_1]$, $x_2 \in S_2 \subset I_2 = (0, M_2]$. Other domain limits can be established in this subsets, such as $x_1 > x_2$.
- $\tilde{\beta} = (\tilde{\beta}_1, \tilde{\beta}_2, \dots, \tilde{\beta}_k)$, a triangular fuzzy number vector.
- $\mu(\beta_j)$ is the membership function of $\beta_j \in \tilde{\beta}_j$.
- $\mu_{\beta_j}(\alpha) = \{\beta_j \mid \mu(\beta_j) \geq \alpha, \alpha \in (0, 1)\}$ set called α -cut.

These sets are closed and bounded, so that is possible to define, for a fuzzy number $\tilde{\beta}_j : \tilde{\beta}_j[\alpha] = [b_1(\alpha), b_2(\alpha)]$, where:

- $b_1(\alpha)$ is the lower value β_j in which $\mu(\beta_j) \geq \alpha$, $\beta_j \in \tilde{\beta}_j$.
 - $b_2(\alpha)$ is the higher value β_j in which $\mu(\beta_j) \geq \alpha$, $\beta_j \in \tilde{\beta}_j$.
 - and $\tilde{\beta}[\alpha] = \prod_j \tilde{\beta}_j[\alpha]$
 - $\tilde{V}(x_1, x_2, \tilde{\beta})$ is a positive and continuous function in $(x_1, x_2) \in S_1 \times S_2$ with partials D_{x_1}, D_{x_2} . This function must be also strictly increasing or strictly decreasing in $x_2 \in S_2$, that is $\tilde{V}(k, x_2, \tilde{\beta})$ is strictly increasing or strictly decreasing for all constant $k \in \mathbb{R}$.
- The fuzzy character of $\tilde{V}(x_1, x_2, \tilde{\beta})$ shown by the tilde placed over V is fixed by $\tilde{\beta}$, and support the use of Buckley-Feuring derivative for solving FPDE.

Date: September 10, 2008.

1991 *Mathematics Subject Classification.* 03E72, 46S40.

Key words and phrases. Fuzzy differential equations, Buckley-Feuring solution, non-polynomial.

- $\varphi(D_{x_1}, D_{x_2})$ is an expression with constant coefficients in (D_{x_1}, D_{x_2}) applied to $\tilde{V}(x_1, x_2, \tilde{\beta})$.
- $F(x_1, x_2, \tilde{\beta})$ continuous function in $(x_1, x_2) \in S_1 \times S_2$.

The specific FPDE treated in this paper has the following form according with this notation:

$$\varphi(D_{x_1}, D_{x_2})\tilde{V}(x_1, x_2, \tilde{\beta}) = \frac{\partial \tilde{V}/\partial x_1}{\partial \tilde{V}/\partial x_2} = F(x_1, x_2, \tilde{\beta})$$

1. THE BUCKLEY-FEURING (B-F) SOLUTION

The Buckley-Feuring (B-F) solution uses a solution of the crisp partial differential equation:

$$V(x_1, x_2) = G(x_1, x_2, \beta),$$

with G continuous $\forall (x_1, x_2) \in S_1 \times S_2$.

The next step is the fuzzification of G :

$$\tilde{Y}(x_1, x_2) = \tilde{G}(x_1, x_2, \tilde{\beta}),$$

with $\tilde{G}$ continuous $\forall (x_1, x_2) \in S_1 \times S_2$ and strictly monotone for $x_2 \in S_2$. Note that $\tilde{Y}_i$ is only the fuzzy representation of G , but not necessary the solution of the fuzzy partial differential equation. If it finally happens and $\tilde{Y}(x_1, x_2)$ is a B-F solution, then $\tilde{V}(x_1, x_2, \tilde{\beta}) = \tilde{Y}(x_1, x_2)$.

With this notation, it is possible to see that:

$$\tilde{Y}(x_1, x_2)[\alpha] = [y_1(x_1, x_2, \alpha), y_2(x_1, x_2, \alpha)], \text{ and}$$

$$\tilde{F}(x_1, x_2, \tilde{\beta})[\alpha] = [f_1(x_1, x_2, \alpha), f_2(x_1, x_2, \alpha)], \forall \alpha.$$

and, by definition:

$$\begin{aligned} y_1(x_1, x_2, \alpha) &= \min\{G(x_1, x_2, \beta), \quad \beta \in \tilde{\beta}[\alpha]\}, \\ y_2(x_1, x_2, \alpha) &= \max\{G(x_1, x_2, \beta), \quad \beta \in \tilde{\beta}[\alpha]\}, \text{ and} \end{aligned}$$

$$\begin{aligned} f_1(x_1, x_2, \alpha) &= \min\{F(x_1, x_2, \beta), \quad \beta \in \tilde{\beta}[\alpha]\}, \\ f_2(x_1, x_2, \alpha) &= \max\{F(x_1, x_2, \beta), \quad \beta \in \tilde{\beta}[\alpha]\}, \\ &\forall x_1, x_2, \alpha. \end{aligned}$$

If it is possible to apply the $\varphi(D_{x_1}, D_{x_2})$ operator to $y_i, i = 1, 2$, getting continuous expressions $\forall (x_1, x_2) \in S_1 \times S_2, \forall \alpha$, then it will be feasible to define the following expression in this domain $\Gamma(x_1, x_2, \alpha)$:

$$\Gamma(x_1, x_2, \alpha) = [\Gamma_1(x_1, x_2, \alpha), \Gamma_2(x_1, x_2, \alpha)]$$

with:

$$\begin{aligned} \Gamma_1(x_1, x_2, \alpha) &= \varphi(D_{x_1}, D_{x_2})y_1(x_1, x_2, \alpha) \\ \Gamma_2(x_1, x_2, \alpha) &= \varphi(D_{x_1}, D_{x_2})y_2(x_1, x_2, \alpha) \end{aligned}$$

For a B-F solution Y , it is necessary to be a fuzzy number for this one. If, for each pair $(x_1, x_2) \in S_1 \times S_2$, $\Gamma(x_1, x_2, \alpha)$ defines an α -cut of a fuzzy number, then, Buckley and Feuring [2] call to $\tilde{Y}_i(x_1, x_2)$ **differentiable**, and we can write:

$$\varphi(D_{x_1}, D_{x_2})\tilde{Y}(x_1, x_2)[\alpha] = \Gamma(x_1, x_2, \alpha),$$

$$\forall (x_1, x_2) \in S_1 \times S_2, \quad \forall \alpha.$$

So that, it is necessary to test if $\Gamma(x_1, x_2, \alpha)$ really define an α -cut of a fuzzy number and verify the differentiability of $\tilde{Y}(x_1, x_2)$. For a triangular fuzzy number, the conditions are [4] :

- (1) $\varphi(D_{x_1}, D_{x_2})y_1(x_1, x_2, \alpha)$ is an increasing function of α , for each $(x_1, x_2) \in S_1 \times S_2$.
- (2) $\varphi(D_{x_1}, D_{x_2})y_2(x_1, x_2, \alpha)$ is a decreasing function of α , for each $(x_1, x_2) \in S_1 \times S_2$.

$$(3) \quad \varphi(D_{x_1}, D_{x_2})y_1(x_1, x_2, 1) \leq \varphi(D_{x_1}, D_{x_2})y_2(x_1, x_2, 1) \text{ for each } (x_1, x_2) \in S_1 \times S_2.$$

Once delimited the differentiability concept of $\tilde{Y}(x_1, x_2)$, it is possible to define the Buckley-Feuring solution. $\tilde{Y}_i(x_1, x_2)$ is a Buckley-Feuring solution if the following conditions hold:

- (1) $\tilde{Y}_i(x_1, x_2)$ is differentiable.
- (2) $\varphi(D_{x_1}, D_{x_2})\tilde{Y}(x_1, x_2) = \tilde{F}(x_1, x_2, \tilde{\beta})$.

Obviously, if differentiability conditions hold by the candidate to B-F solution $\tilde{Y}(x_1, x_2)$, this one will be a fuzzy number. To complete the conditions for B-F solutions only is necessary to test that:

$$\varphi(D_{x_1}, D_{x_2})\tilde{Y}(x_1, x_2) = \tilde{F}(x_1, x_2, \tilde{\beta}),$$

or the equivalent condition:

- (1) $\varphi(D_{x_1}, D_{x_2})y_1(x_1, x_2, \alpha) = F_1(x_1, x_2, \alpha)$.
- (2) $\varphi(D_{x_1}, D_{x_2})y_2(x_1, x_2, \alpha) = F_2(x_1, x_2, \alpha)$.

$$\forall (x_1, x_2) \in S_1 \times S_2, \quad \forall \alpha.$$

In this case we can identify $\tilde{Y}(x_1, x_2)$ with $\tilde{V}(x_1, x_2, \tilde{\beta})$.

2. BOUNDARY CONDITIONS

The FPDE can be subject to certain boundary conditions in a big variety of forms depending on a constant vector $\mathbf{c} = (c_1, \dots, c_n) \in C_1 \times \dots \times C_n$. The inclusion of this ones has not great consequences in the methodology exposed.

The crisp solution G acquires under boundary conditions the form $G(x_1, x_2, \beta, \mathbf{c})$. The fuzzification of G can take $\mathbf{c}$ in a triangular fuzzy vector $\tilde{\mathbf{c}} = (\tilde{c}_1, \dots, \tilde{c}_n) \in C_1 \times \dots \times C_n$ with $\tilde{\mathbf{c}}[\alpha] = \prod_i \tilde{C}_i[\alpha]$, $i = 1, \dots, n$ and:

$$\tilde{Y}(x_1, x_2) = \tilde{G}(x_1, x_2, \tilde{\beta}, \tilde{\mathbf{c}})$$

In this environment with boundary conditions, it is necessary to add a new condition for a B-F solution: $\tilde{Y}(x_1, x_2)$ must satisfy these conditions. In this form $\tilde{Y}_i(x_1, x_2)$ is a Buckley-Feuring solution if the following conditions hold:

- (1) $\tilde{Y}_i(x_1, x_2)$ is differentiable.
- (2) $\varphi(D_{x_1}, D_{x_2})\tilde{Y}(x_1, x_2) = \tilde{F}(x_1, x_2, \tilde{\beta})$.
- (3) $\tilde{Y}(x_1, x_2)$ satisfies the boundary conditions.

3. EXAMPLE

The following FPDE is proposed:

$$\frac{\partial \tilde{V} / \partial x_1}{\partial \tilde{V} / \partial x_2} = \tilde{\beta} x_1^{-1} x_2, \quad \tilde{\beta} \in (0, 1), x_1 \geq 1, x_2 > 0$$

In this example:

$$\tilde{F}(x_1, x_2, \tilde{\beta}) = \tilde{\beta} x_1^{-1} x_2, \quad \tilde{\beta} \in (0, 1), x_1 \geq 1, x_2 > 0$$

and, by definition, the operator

$$\varphi(D_{x_1}, D_{x_2})V(x_1, x_2) \longrightarrow \frac{\partial \tilde{V} / \partial x_1}{\partial \tilde{V} / \partial x_2},$$

A possible solution to this FPDE in a crisp environment is, without special boundary conditions:

$$G(x_1, x_2, \beta) = x_1^\beta x_2 + \gamma, \beta \in (0, 1), x_1 \geq 1, x_2 > 0$$

with $\beta = (\beta, \gamma)$

Applying the fuzzification in β and γ , these ones acquire a triangular fuzzy number form and $\tilde{\beta} = (\tilde{\beta}, \tilde{\gamma})$, $\tilde{\beta} \in (0, 1)$. While G holds all the conditions required, the Buckley-Feuring solution candidate is:

$$\tilde{Y}(x_1, x_2) = G(x_1, x_2, \tilde{\beta}, \tilde{\gamma}) = x_1^{\tilde{\beta}} x_2 + \tilde{\gamma}, \tilde{\beta} \in (0, 1), x_1 \geq 1, x_2 > 0$$

The fuzzy parameters have a membership function associated $\mu(\beta)$ and $\mu(\gamma)$ respectively. From the α -cuts, it is possible to define:

$$\begin{aligned} \tilde{\beta}[\alpha] &= [b_1(\alpha), b_2(\alpha)], \\ \tilde{\gamma}[\alpha] &= [g_1(\alpha), g_2(\alpha)], \text{ and} \\ \tilde{\beta}[\alpha] &= \tilde{\beta}[\alpha] \times \tilde{\gamma}[\alpha] \end{aligned}$$

And from this ones:

$$\begin{aligned} \tilde{Y}(x_1, x_2)[\alpha] &= [y_1(x_1, x_2, \alpha), y_2(x_1, x_2, \alpha)], \text{ and} \\ \tilde{F}(x_1, x_2, \tilde{\beta})[\alpha] &= [f_1(x_1, x_2, \alpha), f_2(x_1, x_2, \alpha)], \forall \alpha. \end{aligned}$$

Where:

$$\begin{aligned} y_1(x_1, x_2, \alpha) &= \min\{G(x_1, x_2, \beta), \beta \in \tilde{\beta}[\alpha]\} = G(x_1, x_2, g_1(\alpha), b_1(\alpha)), \\ y_2(x_1, x_2, \alpha) &= \max\{G(x_1, x_2, \beta), \beta \in \tilde{\beta}[\alpha]\} = G(x_1, x_2, g_2(\alpha), b_2(\alpha)) \end{aligned}$$

and,

$$\begin{aligned} f_1(x_1, x_2, \alpha) &= \min\{F(x_1, x_2, \beta), \beta \in \tilde{\beta}[\alpha]\} = F(x_1, x_2, g_1(\alpha), b_1(\alpha)), \\ f_2(x_1, x_2, \alpha) &= \max\{F(x_1, x_2, \beta), \beta \in \tilde{\beta}[\alpha]\} = F(x_1, x_2, g_2(\alpha), b_2(\alpha)), \end{aligned}$$

$$\forall \alpha \quad x_1 \geq 1, x_2 > 0$$

In the $\tilde{G}$ function proposed, $\forall (x_1, x_2) x_1 \geq 1, x_2 > 0$:

$$\begin{aligned} y_1(x_1, x_2, \alpha) &= x_1^{b_1(\alpha)} x_2 + g_1(\alpha), \quad b_1(\alpha) \in (0, 1) \\ y_2(x_1, x_2, \alpha) &= x_1^{b_2(\alpha)} x_2 + g_2(\alpha), \quad b_2(\alpha) \in (0, 1) \end{aligned}$$

$$\begin{aligned} f_1(x_1, x_2, \alpha) &= b_1(\alpha) x_1^{-1} x_2, \quad b_1(\alpha) \in (0, 1) \\ f_2(x_1, x_2, \alpha) &= b_2(\alpha) x_1^{-1} x_2, \quad b_2(\alpha) \in (0, 1) \end{aligned}$$

Testing the differentiability of $\tilde{G}(x_1, x_2, \tilde{\beta})$, from $\Gamma(x_1, x_2, \alpha) = [\varphi(D_{x_1}, D_{x_2})y_1(x_1, x_2, \alpha), \varphi(D_{x_1}, D_{x_2})y_2(x_1, x_2, \alpha)]$ verifying if $\Gamma(x_1, x_2, \alpha)$ defines an α -cut of a triangular fuzzy number for each pair $(x_1, x_2), x_1 \geq 1, x_2 > 0$, the following conditions must be hold:

- (1) $\varphi(D_{x_1}, D_{x_2})y_1(x_1, x_2, \alpha) = b_1(\alpha)x_1^{-1}x_2$, $b_1(\alpha) \in (0, 1)$ is an increasing function of α , for each pair $(x_1, x_2), x_1 \geq 1, x_2 > 0$.

It will happen if $\varphi(D_{x_1}, D_{x_2})y_1(x_1, x_2, \alpha)$ has positive derivative on α :

$$d_\alpha (b_1(\alpha)x_1^{-1}x_2) = b_1(\alpha)'x_1^{-1}x_2$$

While β is a triangular fuzzy number and $b_1(\alpha)$ is defined from its α -cuts, $b_1(\alpha)$ satisfies the condition and is an increasing function with $b_1(\alpha)' > 0$, and $x_1^{-1}x_2 > 0$ for $x_1 \geq 1, x_2 > 0$, $d_\alpha (b_1(\alpha)x_1^{-1}x_2) > 0$ and the condition holds.

- (2) $\varphi(D_{x_1}, D_{x_2})y_2(x_1, x_2, \alpha) = b_2(\alpha)x_1^{-1}x_2$, $b_2(\alpha) \in (0, 1)$ is a decreasing function of α , for each pair $(x_1, x_2), x_1 \geq 1, x_2 > 0$.

Again, it will happen if $\varphi(D_{x_1}, D_{x_2})y_2(x_1, x_2, \alpha)$ has negative derivative on α :

$$d_\alpha (b_2(\alpha)x_1^{-1}x_2) = b_2(\alpha)'x_1^{-1}x_2$$

We can now use that β is a triangular fuzzy number and $b_2(\alpha)$ is defined from its α -cuts, so that, in analogy, $b_2(\alpha)$ satisfies the condition and is a decreasing function with $b_2(\alpha)' < 0$, and $x_1^{-1}x_2 > 0$ for $x_1 \geq 1, x_2 > 0$, $d_\alpha(b_1(\alpha)x_1^{-1}x_2) < 0$ and the condition holds.

$$(3) \quad \varphi(D_{x_1}, D_{x_2})y_1(x_1, x_2, 1) \leq \varphi(D_{x_1}, D_{x_2})y_2(x_1, x_2, 1) \forall x_1, x_2 \in I_1 \times I_2.$$

In the example:

$$b_1(1)x_1^{-1}x_2 \leq b_2(2)x_1^{-1}x_2$$

This condition is holds automatically if we realize that $b_1(\alpha)$ and $b_2(\alpha)$ are defined from a triangular fuzzy number and $b_2(1) > b_1(1)$

At this point, it is possible to say that $\tilde{Y}(x_1, x_2) = G(x_1, x_2, \tilde{\beta}, \tilde{\gamma}) = x_1^{\tilde{\beta}}x_2 + \tilde{\gamma}, \tilde{\beta} \in (0, 1), x \geq 1$ is differentiable and good candidate for B-F solution. But it is necessary the last step, and it must satisfier that:

$$\varphi(D_{x_1}, D_{x_2})\tilde{Y}(x_1, x_2) = \tilde{F}(x_1, x_2, \tilde{\beta}) \quad \beta, \gamma \in (0, 1), \forall (x_1, x_2), x_1 \geq 1, x_2 > 0.$$

For $G(x_1, x_2, \tilde{\beta}, \tilde{\gamma}) = x_1^{\tilde{\beta}}x_2 + \tilde{\gamma}, \tilde{\beta} \in (0, 1), x \geq 1, x_2 \geq 0$:

$$\varphi(D_{x_1}, D_{x_2})\tilde{Y}(x_1, x_2) = \tilde{\beta}x_1^{-1}x_2, \quad \tilde{\beta} \in (0, 1), x_1 \geq 1, x_2 > 0$$

and,

$$\tilde{F}(x_1, x_2, \tilde{\beta}) = \tilde{\beta}x_1^{-1}x_2, \quad \tilde{\beta} \in (0, 1), x_1 \geq 1, x_2 > 0.$$

Thus $G(x_1, x_2, \tilde{\beta}, \tilde{\gamma}) = x_1^{\tilde{\beta}}x_2 + \tilde{\gamma}, \tilde{\beta} \in (0, 1), x \geq 1, x_2 \geq 0$ is a Buckley-Feuring solution for this non-polynomial fuzzy partial differential equation.

REFERENCES

- [1] Allahviranloo, T. Diference methods for fuzzy partial differential equations, Computational Methods in Applied Mathematics, Vol.2 (n°3) (2002), pp. 233-242.
- [2] Buckley, J.J. and Feuring, T. Introduction to fuzzy partial differential equations, Fuzzy Sets and Systems, 105 (1999), pp.241-248.
- [3] Buckley, J.J. and Feuring, T. Fuzzy differential equations, Fuzzy Sets and Systems, 110 (2000), pp.43-54.
- [4] Goetschel, R. and Voxman, W. Elementary fuzzy calculus, Fuzzy Sets and Systems, 18 (1986), pp.319-330.
- [5] Quan, H., Hua, Y. and Jones, J.D. A general method for calculating functions of fuzzy numbers, Applied Mathematics Letters, Vol.5 (n°6) (1992), pp.51-55.
- [6] Seikkala, s. On the fuzzy initial value problem, Fuzzy Sets and Systems, 24 (1987), pp.31-43.
- [7] Zadeh, L.A. Fuzzy Sets. Inf. Control, 8 (1965), pp.338-353.

FACULTAD DE MATEMÁTICAS. UNIVERSIDAD DE SEVILLA, 41012 SEVILLA, SPAIN

Keywords: Fuzzy differential equations, Buckley-Feuring solution, non-polynomial

2000 Mathematics Subject Classification: 03E72, 46S40

E-mail address: davidgalvez@us.es; jlpino@us.es